

A randomized controlled trial of technology-enhanced behavioral parent training: sustained parent skill use and child outcomes at follow-up

Justin Parent,^{1,2,3}  Margaret T. Anton,⁴ Raelyn Loiseau,⁵ April Highlander,⁵ Nicole Breslend,⁶ Rex Forehand,⁶ Megan Hare,³ Jennifer K. Youngstrom,⁵ and Deborah J. Jones⁵

¹Warren Alpert Medical School, Brown University, Providence, RI, USA; ²Bradley/Hasbro Children's Research Center, E. P. Bradley Hospital, Providence, RI, USA; ³Center for Children and Families, Florida International University, Miami, FL, USA; ⁴AbleTo Inc., New York, NY, USA; ⁵University of North Carolina at Chapel Hill, Chapel Hill, NC, USA; ⁶University of Vermont, Burlington, VT, USA

Background: Early-onset (3–8 years old) disruptive behavior disorders (DBDs) have been linked to a range of psychosocial sequelae in adolescence and beyond, including delinquency, depression, and substance use. Given that low-income families are overrepresented in statistics on early-onset DBDs, prevention and early-intervention targeting this population is a public health imperative. The efficacy of Behavioral Parent Training (BPT) programs such as Helping the Noncompliant Child (HNC) has been called robust; however, given the additional societal and structural barriers faced by low-income families, family engagement and retention barriers can cause effects to wane with time. This study extends preliminary work by examining the potential for a Technology-Enhanced HNC (TE-HNC) program to improve and sustain parent skill proficiency and child outcomes among low-income families.

Methods: A randomized controlled trial with two parallel arms was the design for this study. A total of 101 children (3–8-years-old) with clinically significant problem behaviors from low-income households were randomized to HNC ($n = 54$) or TE-HNC ($n = 47$). Participants were assessed at pre-treatment, post-treatment, 3-month, and 6-month follow-ups. Primary outcomes were parent-reported and observed child behavior problems. Secondary outcomes included observed parenting skills use (ClinicalTrials.gov Identifier: NCT02191956). **Results:** Primary analyses used latent curve modeling to examine treatment differences in the trajectory of change during treatment, maintenance of treatment gains, and levels of outcomes at the 6-month follow-up. Both programs yielded improvements in parenting skills and child problems at post-treatment. However, TE-HNC families evidenced greater maintenance of parent-reported and observed child behavior and observed positive parenting skills at the 6-month follow-up. **Conclusions:** Our findings contribute to an ongoing line of work suggesting that technology-enhanced treatment models hold promise for increasing markers of engagement in BPT and sustaining long-term outcomes among low-income families. **Keywords:** Behavioral parent training; technology; low-income families; child behavior disorders.

Introduction

Early-onset disruptive behavior disorders (DBDs; oppositional defiant disorder, conduct disorder) co-occurring with attention-deficit/hyperactivity disorder are the second-leading cause of child mental health referrals worldwide (Merikangas, Nakamura, & Kessler, 2009). Early-onset DBDs affect an estimated 113 million youth (Polanczyk, Salum, Sugaya, Caye, & Rohde, 2015), increasing their vulnerability for later delinquency and antisocial behavior, depression, substance use, academic underachievement, relationship and employment instability, and chronic illness (e.g. Fergusson, Horwood, & Ridder, 2005; Odgers et al., 2007; Piquero et al., 2016). Therefore, early identification and treatment of DBDs are critical, with estimates suggesting these

efforts can save \$2.6 to \$4.4 million per high-risk child (Cohen & Piquero, 2009).

Behavioral Parent Training (BPT), a family of evidence-based treatments with a common history, theory, and practice elements, is the recommended first-line treatment for children with DBDs (see Comer, Chow, Chan, Cooper-Vince, & Wilson, 2013; Kaminski & Claussen, 2017 for reviews). Research suggests that BPT, which targets the coercive cycle of parent-child interaction implicated in the etiology and maintenance of early-onset DBDs, is more effective than any other treatment approach to date (see Chorpita et al., 2011; Leijten, Raaijmakers, de Castro, & Matthys, 2013 for reviews). Indeed, the impact of BPT has been called relatively “robust,” yielding moderate effects for improvements in parenting ($d = .45$) and child behavior ($d = .42$) (see Lundahl, Risser, & Lovejoy, 2006, for a review). Moreover, data suggest that BPT efficacy is equivalent regardless of socioeconomic status and particularly when the child's behavior is clinically significant (see Leijten et al., 2013, for a

Conflict of interest statement: No conflicts declared.

Ethical considerations: Written consent from the subjects was obtained.

meta-analysis). However, BPT effects wane after treatment, likely due to a number of factors including reduction in parent use of skills in the context of families' daily lives (see Lundahl et al., 2006 for a review).

Maintaining skill use after treatment may be particularly challenging for low-income families, given that financial strain and related stressors are known to increase risk for the coercive cycle and vulnerability for child problem behavior (see Masarik & Conger, 2017; Shaw & Shelleby, 2014 for reviews). Further, evidence suggests that stressors related to economic disadvantage may be confounded or exacerbated by families' sociocultural experiences related to their race and ethnicity (e.g. racism, discrimination), and in turn, differentially impact families' ability to experience positive outcomes when engaging in BPT. Indeed, studies suggest that in comparison to White, non-Latinx populations, racial and ethnic minority families may not have the opportunity to equally benefit from BPT in part, due to their disproportionate experiences of socioeconomic disadvantage and parental stress, contributing to lower engagement and completion rates (Lavigne et al., 2010; McCabe, Yeh, & Zerr, 2020). Additionally, financial strain also likely competes with caregivers' ability to effectively use BPT skills after treatment completion when faced with disproportionate adversity and daily stressors (see Jones et al., 2013; Lundahl et al., 2006 for reviews).

Consistent with efforts to increase in- and out-of-session support to low-income families throughout BPT (e.g. Chacko, Wymbs, Chimiklis, Wymbs, & Pelham, 2012; Williams, Lynch, & Glasgow, 2007), the field must also consider how to help low-income families maintain treatment gains at follow-ups. One such approach capitalizes on the rise of telehealth generally and the uptake of mobile phones among low-income consumers in particular (e.g. Georgeson, Highlander, Loiselle, Zachary, & Jones, 2020; Jones et al., 2013; Lindhiem, Bennett, Rosen, & Silk, 2015). Telehealth broadly refers to the use of technology as an adjunct to or replacement for standard health services with the goal of increasing service accessibility, engagement, and/or outcomes. Although telehealth has received increased focus due to the COVID-19 pandemic and associated social distancing mandates, technology ranging from the relatively early and basic (e.g. videotape modeling) to those more recent and sophisticated (e.g. remote coaching, internet-delivered sessions) has long been rooted in the literature on BPT (e.g. Chacko, Isham, Cleek, & McKay, 2016; Jones et al., 2013; Ortiz, Vidair, Acri, Chacko, & Kobak, 2020, for a review). Consistent with national trends, low-income homes are cutting the cord on landlines; however, they are more likely than high-income families to rely on mobile phones as the primary and often only digital device in the home (e.g. Blumberg & Luke, 2018; Pew Research Center,

2019). This trend makes sense, given that mobile phones can cost-effectively increase low-income parents' ability to use a range of features (e.g. text/email, electronic/shared calendars) essential to parenting children in the 21st century.

Building upon these trends, this study examined parent skill use and child behavior at post-treatment, as well as 3- and 6-month follow-up, in 101 low-income families randomized to *Helping the Noncompliant Child* (HNC; McMahon & Forehand, 2003) or to Technology-Enhanced HNC (TE-HNC). To date, results across two TE-HNC trials suggest the potential to cost-effectively and efficiently (i.e. less time to master program skills) improve aspects of low-income families' engagement in BPT without compromising parent satisfaction with services (Jones et al., 2014, 2021). Effect sizes in preliminary work also suggested greater TE-HNC improvement in child behavior at post-treatment (Jones et al., 2014). This study extended this pilot with a larger sample, long-term follow-up, and more rigorous statistical methods to evaluate post-treatment and follow-up effects for parent skill use and child behavior in this larger randomized controlled trial. We hypothesized that technology enhancements (TE-HNC) would help families maintain treatment gains for parenting and child behaviors resulting in superior long-term outcomes as compared to standard care as usual (HNC).

Methods

Participants

This study included 101 children (3–8-years-old) from families with low-income (<250% of the Federal Poverty Guidelines) and their primary caregivers (see Table 1). The majority of parents identified as White (68.3%) or Black/African American (21.8%) with an additional 1.0% as Native American and 8.9% identifying as mixed race. Further, 6.9% of parents identified as Hispanic/Latinx. Similarly, most children were White (63.4%) or Black/African American (20.8%), with 1.0% identifying as Native American and 13.9% identifying as mixed race. Further, 13.9% of children were Hispanic/Latinx. Children with clinically significant problem behaviors (*Eyberg Child Behavior Inventory Problem* > 15 or *Intensity* > 131; Eyberg & Pincus, 1999) were included. Children were excluded if they had a significant developmental and/or physical impairment that prohibited use of HNC. Families were also excluded if caregivers had a current mood, psychotic, and/or substance use disorder or a pending and/or prior substantiated child abuse/neglect case. Participants were recruited via advertisements and flyers distributed at nonprofit organizations, local schools, agencies serving low-income families, and word-of-mouth (see Khavjou et al., 2020; Khavjou, Turner, & Jones, 2018).

Procedure

Families completed a phone screen and baseline assessment at a community-based clinic to confirm eligibility and provide consent for their family's participation. Eligible families were then randomized 1:1 to either HNC or TE-HNC. Families were compensated \$50 per assessment for completing the baseline, post-, 3-, and 6-month follow-up assessments. All procedures were approved by the university's institutional review board.

Table 1 Demographics by group at baseline

	Total Sample (<i>N</i> = 101) <i>M</i> (<i>SD</i>)	HNC (<i>n</i> = 54) <i>M</i> (<i>SD</i>)	TE-HNC (<i>n</i> = 47) <i>M</i> (<i>SD</i>)
Child			
Age (years)	4.19 (1.19)	4.28 (1.17)	4.13 (1.19)
Gender (% male)	54.9%	57.41%	53.19%
Race %			
White	63.4%	57.41%	70.21%
Black/African American	20.8%	25.93%	14.89%
Native American/Alaskan	1.0%	0.00%	2.13%
Multiracial	13.9%	14.81%	12.77%
Not reported	1.0%	1.85%	0.00%
Hispanic/Latinx	13.9%	14.81%	12.77%
Parent			
Age (years)	31.66 (6.72)	32.50 (6.12)	31.34 (5.88)
Gender (% female)	97.06%	98.15%	97.87%
Race %			
White	68.3%	62.96%	74.47%
Black/African American	21.8%	27.78%	14.89%
Native American/Alaskan	1.0%	0.00%	2.13%
Multiracial	7.9%	7.41%	8.51%
Not reported	1.0%	1.85%	0.00%
Hispanic/Latinx	6.9%	7.41%	6.38%
Marital status			
Single	24.51%	27.78%	21.28%
Married/living together	61.76%	59.26%	63.83%
Divorced/separated	13.73%	12.96%	14.80%
Employed in Any capacity	46.08%	50.00%	40.43
Smartphone owner	94.1%	92.6%	95.7%
iPhone	44.2%	38.0%	51.1%
Android	54.7%	62.0%	46.7%
Other	1.1%	0%	2.2%

Intervention

All families received HNC, which is a therapist-delivered, criteria-based (i.e. therapists conduct weekly observation and coding of skill use to determine progression through skills and program completion) BPT intervention for children with behavior disorders. HNC included weekly face-to-face therapy sessions (60 min), as well as a brief midweek phone check-in. HNC consists of two phases: Differential Attention (e.g. increasing positive attention, ignoring inappropriate behavior) and Compliance Training (e.g. utilizing time outs). When parents progress to Phase II (i.e. Compliance Training), they continue to practice Phase I skills to maintain skill proficiency.

Families in the TE-HNC group received the full HNC protocol (McMahon & Forehand, 2003) augmented with a digital companion, Tantrum Tamers©. Tantrum Tamers is a HIPAA-compliant, interactive system that allowed therapists to monitor caregiver activity on the mobile application, as well as tailor the focus and pace of treatment based on parent practice and progress. The Tantrum Tamers application included: (a) daily surveys of skills practice, (b) weekly video-recorded home practice, (c) daily text reminders for skill practice and appointments; (d) video calls with the family midweek to problem solve obstacles; and (e) skills video series to model new skills and

share with other caregivers (see Jones et al., 2021 for more information). Additionally, a homework checklist was added to remind caregivers of daily and weekly assignments. TE-HNC families had access to all app functionality during treatment. After families completed treatment, they had access to a limited range of content, including the skills video series and surveys with automated feedback through their 3-month follow-up assessment when they returned their study phones. From 3- to 6-month follow-ups, TE-HNC parents had access to a program blog, which provided content intended to remind caregivers of program content.

Therapist training and fidelity

Master's-level therapists treated families in both groups. Training included reviewing treatment manuals, establishing reliability with the HNC coding system, role-play and session observations and discussion, weekly observation, and supervision and feedback by two licensed clinical psychologists. Approximately, a quarter of sessions (24%) were coded by one Master's-level coder to ensure treatment fidelity (97% fidelity); 72% of those were double-coded (90% reliability between coders). In addition, 35% of sessions were coded for therapist competence by at least one doctoral-level coder and of those 22% were coded by a second doctoral-level coder, yielding an average competence rating of 97%.

Measures

Demographics. Caregivers reported their and their child's demographic information at baseline, including, age, race/ethnicity, marital status, education level, and income.

Parent-reported child problem behavior. At all waves, caregivers completed the *Eyberg Child Behavior Inventory* (ECBI; Eyberg & Robinson, 1983). The ECBI is a 36-item caregiver-report inventory measuring common disruptive behavior problems in youth ages 2–16. The ECBI has two scales: (1) the Intensity Scale, which measures the frequency ($1 = \text{never}$, $7 = \text{always}$) with which the child engages in each of the 36 behaviors (Range 36–252; ≥ 131 clinically significant), and (2) the Problem Scale, which asks caregivers to indicate whether each of the 36 behaviors is “a problem for you” (*yes* or *no*; Range 0–36; ≥ 15 clinically significant). The ECBI is sensitive to BPT interventions (e.g. Nixon, Sweeney, Erickson, & Touyz, 2003; Webster-Stratton & Hammond, 1997) and has demonstrated internal consistency and test-retest reliability (Eyberg & Robinson, 1983; Funderburk, Eyberg, Rich, & Behar, 2003). In this study, the omega reliability coefficients ranged from .88 to .93 for the Intensity Scale and .82 to .92 for the Problem Scale across waves.

Observed parenting and child compliance. Parent-child observations were conducted at all waves. Coders received approximately 50 hr of training in the Behavioral Observation Coding System (McMahon & Forehand, 2003) and reached at least 80% agreement on one or more of the coded behaviors with expert coders on a series of training videos. Half of the videos were double-coded for fidelity. When two coders failed to reach 80% agreement, they jointly code the observation to resolve discrepancies. Behaviors are reported at a rate per minute during a 5-min observation period to account for variability in interaction length. *Attends* was defined as positive attention in which the parent provides an ongoing verbal description of what the child is doing and *Rewards* was defined as positive attention that is provided following the child's appropriate behavior, and these were combined for a single average score. *Questions* (an interrogation to which the only appropriate response is verbal) and *Instructions* (parent-issued

command toward the child), were also measured by the average number of caregiver Questions and Instructions per minute. Finally, child *Compliance* was measured as the percentage of all parent Clear Instructions the child complied with within 5 s after a command was issued.

Data analytic plan

Managing missing data and intent-to-treat analysis approach. Prior to conducting analyses, the pattern of missingness across follow-ups was examined to determine if data were missing at random. Multiple imputation was conducted for use with descriptive and group mean statistics. Twenty imputed datasets were created by group using Blimp 2.2.3 software (Keller & Enders, 2019) using a fully conditional specification and Gibbs sampler Markov chain Monte Carlo (MCMC) algorithm. MI diagnostics were conducted to ensure that the potential scale reduction values were less than 1.05 (Gelman & Rubin, 1992). For primary group difference analyses, full information maximum likelihood (FIML) estimation techniques were used for inclusion of all available data based on intent-to-treat guidelines.

Treatment effect sizes. To examine effect sizes, Cohen's *d* was used to compare the magnitude of the experimental effects within each treatment and across groups (Cohen, 1988). Within-group effect sizes were used to examine magnitude in change in child problem behaviors and observed parenting comparing pre-treatment and post-treatment, pre-3-month follow-up, and pre-6-month follow-up, controlling for the correlation between each time point, for each group. To test the clinical significance of treatment effects, we utilized the Reliable Change Index (RCI; Jacobson & Truax, 1991), which suggests that effects attributable to treatment are most likely not due to chance, with a value of 1.96 or greater indicating the effect is reliable.

Comparing treatment conditions. For primary analyses, we employed latent curve modeling (Preacher, Wichman, MacCallum, & Briggs, 2008) to examine differences in the trajectory of change during treatment, maintenance of gains, and outcomes at the 6-month follow-up. Given expected nonlinear patterns of change, we used a spline growth model with a treatment slope (pre- to post-treatment), a maintenance slope (from post through 3- and 6-month follow-ups), and intercept coded as the mean level at the 6-month follow-up. Growth curve analyses were conducted using Mplus 8.2 software (Muthén & Muthén, 2017) and ML estimation with robust standard errors. FIML techniques were used for inclusion of all available data. Primary outcomes were child behavior problems and secondary outcomes were parenting skills.

Results

Figure 1 depicts the CONSORT diagram including participant retention at each follow-up by group. No significant condition differences were observed across demographics. Overall, rates of retention by group were similar across follow-ups. Patterns of missingness did not significantly differ by the following: treatment condition; child age, sex, race/ethnicity; caregiver age, ethnicity/race; nor family economic stress, all *ps* > .05. Further, random patterns of missingness along with a nonsignificant Little's missing completely at random (MCAR) test, $\chi^2(239) = 267.64$, *p* > .05, suggest that the mechanism

of missingness was MCAR and support use of multiple imputation and FIML for primary analyses.

Treatment effect sizes

Child problem behavior. HNC and TE-HNC both demonstrated large within-group effects sizes from pre-post-treatment, pre-3-month follow-up, and pre-6-month follow-up (see Table 2). Further, HNC and TE-HNC both demonstrated clinically significant treatment effects (RCIs > 1.96), indicating improvements.

Parenting and child compliance. Examining within-group effects, both treatments demonstrated large effect sizes from pre-post-treatment, pre-3-month follow-up, and pre-6-month follow-up for observed Questions + Instructions and Attends + Rewards. Results indicate that both treatments were successful in reducing the number of questions and instructions, while improving parents' attending and rewards at the end of treatment with sustained effects over 6 months. Both treatments also improved children's observed compliance.

Treatment comparisons

Parent-reported child problem behavior. Parameter estimates for each outcome are shown in Table 3. Model fit for the ECBI Intensity, $\chi^2(5) = 2.02$, *p* = .846, RMSEA = .000 [90% CI .000, .078], CFI = 1, SRMR = .053, and Problem, $\chi^2(5) = 7.86$, *p* = .175, RMSEA = .073 [.000, .169], CFI = .971, SRMR = .074, models had adequate to excellent fit. As is depicted in Figure 2A, the frequency of parent-reported behavior problems (i.e. intensity scale) demonstrated similar rates of change during the active treatment phase across conditions. However, TE-HNC had superior maintenance of gains resulting in statistically significant group differences at the 6-month follow-up for ECBI Intensity. Further supporting this difference, 85.10% TE-HNC participants were below the clinical cut-off compared to 62.96% of those in HNC at the 6-month follow-up (see Table 2). Results for the ECBI Problem scale (Figure 2B) showed a similar pattern of results, but with group differences only approaching statistical significance. TE-HNC had 10% more children within the normative range, at the 6-month follow-up, compared to HNC. Interpretation of this group difference supports a clinically meaningful difference between conditions for the ECBI Problems outcome.

Observed parenting and child compliance. Parameter estimates for each model are shown in Table 3. Model fit for observed Attends + Rewards, $\chi^2(5) = 5.21$, *p* = .391, RMSEA = .021 [.000, .145], CFI = .995, SRMR = .081, and Questions + Instructions, $\chi^2(5) = 8.91$, *p* = .113, RMSEA = .090 [.000, .185], CFI = .860,




Figure 1 CONSORT diagram

SRMR = .079, demonstrated excellent to adequate fit. Regarding child compliance, we fixed the residual variance of maintenance slope to zero due to a negative residual variance in the initial model. After this adjustment, the model fit was adequate and without errors, $\chi^2(7) = 7.34, p = .395, RMSEA = .023 [.000, .131], CFI = .986, SRMR = .091$. Results depicted in Figure 2C support group differences on observed compliance with a statistically significant difference between conditions. TE-HNC demonstrated 20% higher rates of observed compliance compared to HNC. Regarding observed Attends + Rewards (see Figure 2D), a similar pattern of change emerged in initial pre-to-post improvements in parenting, with improved maintenance of positive parenting by TE-HNC (see Table 3). Specifically, TE-HNC showed approximately 50% greater rates of observed positive parenting behaviors during a structured task at the 6-month follow-up. Lastly, Questions + Instructions

showed similar rates of initial treatment change and maintenance of gains across groups.

Discussion

This study examined the capacity for TE-HNC to improve parenting skills and child behavior at post-treatment and follow-up relative to standard HNC. Consistent with our pilot, effect sizes suggest families in both groups experience clinically significant gains in parenting skills and child behavior. Between group effect sizes favored TE-HNC, which also resulted in a greater percentage of families below clinical cut-offs. However, primary analyses suggested technology enhancements had similar initial treatment gains as standard care with differences beginning to emerge in the long-term maintenance phase and becoming statistically significant at the 6-month follow-up. Specifically, at the 6-month follow-up, TE-HNC demonstrated lower levels of parent-

Table 2 Means, SDs, and effect sizes for parent and child outcomes

	Baseline <i>M</i> (<i>SD</i>)	Post <i>M</i> (<i>SD</i>)	3-Month <i>M</i> (<i>SD</i>)	6-Month <i>M</i> (<i>SD</i>)	Within group			Between group		
					Pre- post	Pre-3 Month	Pre-6 Month	Post Cohen's <i>d</i>	3-Month	6-Month
ECBI intensity (P)										
HNC	155.4 (27.9)	110.25 (23.6)	115.6 (26.9)	118.35 (35.39)	1.74	1.44	1.16	.27	.56	.48
TE-HNC	150.7 (29.9)	103.84 (23.7)	101.3 (24.2)	102.37 (30.10)	1.72	1.80	1.60			
% below clinical cutoff										
HNC	22.22%	77.78%	70.37%	62.96%						
TE-HNC	23.40%	89.36%	89.36%	85.10%						
ECBI problem (P)										
HNC	23.52 (6.47)	13.65 (6.6)	13.29 (9.6)	13.72 (10.8)	1.50	1.24	1.09	.17	.41	.37
TE-HNC	22.48 (5.76)	12.39 (7.9)	9.72 (7.6)	10.01 (9.07)	1.42	1.85	1.60			
% below clinical cutoff										
HNC	3.70%	50.76%	57.03%	56.01%						
TE-HNC	4.47%	68.81%	76.92%	67.66%						
Questions + Instructions										
HNC	3.97 (2.01)	0.78 (0.98)	1.17 (1.58)	1.05 (2.38)	2.00	1.54	1.32	.23	.44	.23
TE-HNC	3.85 (2.00)	0.56 (0.87)	0.58 (0.93)	0.59 (1.49)	2.12	2.08	1.83			
Attends + Rewards (O)										
HNC	1.09 (.91)	6.69 (3.16)	5.96 (4.43)	4.94 (3.10)	2.39	1.51	1.67	.25	.17	.60
TE-HNC	1.18 (1.19)	7.56 (3.72)	6.67 (3.77)	7.57 (5.50)	2.29	1.95	1.59			
Child Compliance (O)										
HNC	0.34 (0.21)	0.52 (0.26)	0.51 (0.28)	0.43 (0.31)	.76	.68	.34	.30	.45	.59
TE-HNC	0.33 (0.24)	0.60 (0.27)	0.65 (0.34)	0.63 (0.36)	1.05	1.08	.97			

ECBI, Eyberg Child Behavior Inventory; P, parent report; O, observation.

Table 3 Model results for parenting and child outcomes

	ECBI intensity		ECBI problems		Compliance		Attends + Rewards		Questions + Instructions	
	<i>b</i> (SE)	<i>p</i>	<i>b</i> (SE)	<i>p</i>	<i>b</i> (SE)	<i>p</i>	<i>b</i> (SE)	<i>p</i>	<i>b</i> (SE)	<i>p</i>
Condition—6M Intercept	-15.54 (6.59)	.018	-3.58 (2.00)	.074	0.203 (0.063)	.001	2.09 (0.731)	.004	-0.484 (0.324)	.136
Condition—treatment slope	0.575 (8.06)	.943	0.003 (1.97)	.999	0.008 (0.088)	.927	-.10 (0.910)	.913	-0.114 (0.483)	.813
Condition—maintenance slope	-3.72 (2.86)	.194	-0.820 (.822)	.319	0.066 (0.035)	.061	.700 (0.394)	.075	-0.081 (0.159)	.609
6M Mean	117.71 (4.68)	.000	13.14 (1.46)	.000	0.424 (0.041)	.000	4.91 (0.370)	.000	1.14 (0.259)	.000
Treatment slope mean	-47.14 (4.94)	.000	-9.55 (1.39)	.000	0.246 (0.070)	.000	6.49 (0.609)	.000	-3.26 (0.317)	.000
Maintenance mean	3.14 (2.12)	.139	-0.276 (0.673)	.682	-0.052 (0.029)	.074	-.889 (0.24)	.000	0.144 (0.118)	.222

6 M, 6-Month Follow-up; Condition, HNC (coded 0) vs. TE-HNC (coded 1); ECBI, Eyberg Child Behavior Inventory.

reported and observed child disruptive behavior and higher levels of observed positive parenting skills relative to standard HNC.

The pattern observed in this study—similar initial improvements but superior long-term maintenance—may make both empirical and clinical sense. That is, HNC is a criteria-based BPT program, which means parents do not move on with each subsequent skill until they demonstrate proficiency of the previous skill. Further, they do not complete the program until they demonstrate proficiency of all Phase I and II skills. Given that progression through treatment is determined by weekly coding of parent and child behavior, progression, and completion of HNC suggests there has been clinically significant change in

parenting skills and child behavior. Thus, criteria-based programs like HNC may limit significant initial differences across groups at post-treatment based on floor or ceiling effects. TE-HNC caregivers did, however, sustain greater improvements in parenting skills across 3- and 6-month follow-ups at a higher level than HNC families. Specifically, TE-HNC families had higher rates of observed positive parenting and compliance and caregivers reported a lower frequency and intensity of child behavior problems by the 6-month follow-up compared to HNC.

One possible reason for the maintained gains in TE-HNC is that parents experienced increased support from the program and therapist throughout treatment (e.g. tailored feedback via surveys and




Figure 2 Mean plots from growth curve model results of disruptive behavior and parenting outcomes

midweek calls.). For example, TE-HNC participants had better midweek call participation and homework compliance as well as rating the usefulness of the overall program higher than HNC participants (Jones et al., 2021). In addition, TE-HNC parents had ongoing access to program content through the 3-month (i.e. Tantrum Tamers app) and 6-month (skills blog) follow-ups, providing additional, although relatively low-level, support. Thus, if parenting is the primary mechanism of change that allows parents to preserve gains, then ongoing technology-supported practice and use of those skills at home is likely to maintain improvements in child behavior as well. In particular, technology-enhancements may have resulted in a continued and deeper learning of program theory and skills and, in turn, greater maintenance of treatment gains (Anton & Jones, 2019). As low-income families are increasing the use of mobile phones (94.1% in this study), yet still face barriers accessing mental health care (e.g. transportation), such technology enhancements may represent a viable option to increase dissemination and implementation in community settings. Further, as reported in Jones et al. (2021), families in TE-HNC completed the program in fewer weeks

($M = 11.63$) than HNC ($M = 14.15$, $p \leq .05$), which offset any additional program costs. Additionally, this increased efficiency did not come at the expense of consumer satisfaction, which was equivalent across groups. Together, this study results and that of Jones et al. (2021) suggest the promise of TE-HNC as a cost-effective approach that increases engagement and delivery efficiency while also achieving improved maintenance of effects for some parenting and child externalizing outcomes.

As with all research this study has limitations. First, we included 3- and 6-month, but not more distal follow-ups. That said, research suggests that BPT outcomes at 6-months are comparable to 12-months (see de Graaf, Speetjens, Smit, de Wolff, & Tavecchio, 2008 for meta-analysis). In addition, we provided all families in TE-HNC with phones and service plans through the 3-month follow-up, which would not be feasible in front-line service settings. Given data suggesting 44% of low-income users let service plans lapse due to finances (Pew Research Center, 2015), ongoing work in this area will have to consider strategies to increase the sustainability of this approach. It was also true that the vast majority of our families already had their own

mobile phone (94.1%). Therefore, supplying and requiring them to use a program phone may have actually been burdensome. Given that our budget precluded paying for phones and plans beyond the 3-month assessment, we provided families with content via a blog. Although it highlights that effects were maintained even after phones were returned and access to the app ended, it also introduces additional variability between the 3- and 6-month assessments. However, this study did not systematically track post-treatment access to the app (prior to the 3-month) or other content via the blog, which limited our ability to fully understand the mechanisms underlying greater maintenance of treatment effects for the TE-HNC group. Future studies would benefit from systematically tracking both active treatment and post-treatment engagement in technology-based supports to inform further refinements aimed at long-term skills maintenance. For example, Jones et al. (2021) found that an eBook-enhanced BPT program did not result in added benefit for parenting outcomes at post-treatment or follow-up. Thus, it may be that technology enhancements that involve therapist interaction during active treatment may be more effective methods for continued and deeper learning of program theory and skills and, in turn, greater maintenance of treatment gains compared to lighter technology-only enhancements (e.g. skill videos). Finally, future research with large sample sizes should explore the potential differential impact of technology-enhanced BPT on long-term maintenance of effects across caregiver race, ethnicity, and education within low-income populations.

Despite such limitations, this is the first study of its kind to show improved follow-up effects with a technology-enhanced BPT model. Given the common elements characterizing this family of interventions, findings should generalize to other BPT programs as well. In addition, we focused on low-income families who are at-risk for early-onset DBDs, yet experience more obstacles to engaging and completing services. Thus, while low-income families are particularly vulnerable, they are often not the focus of intervention research. Lastly, much policy attention and funding has been devoted to the potential for technology to increase engagement in children's mental health. Empirical research on technology-enhanced treatment models, however, primarily reflects pilot

studies limiting conclusions that can be drawn about the efficacy of and next steps in this approach (see Georgeson et al., 2020 for a review). This study moved beyond earlier pilot studies through a larger sample and long-term follow-ups which enhances confidence in and potential for generalizability of findings.

In summary, our findings contribute to an ongoing line of work suggesting that technology-enhanced treatment models hold promise for increasing markers of engagement in BPT (Jones et al., 2021) and sustaining long-term outcomes. As progress is made in intervention research using artificial intelligence and machine learning (e.g. Cui & Gong, 2018; Timmons et al., 2019), future work must consider if additional just-in-time responsiveness in the context of the families' daily life can further bolster effects. While financial strain is a chronic stressor for low-income families, there is also variability within and between parents in terms of their day-to-day experience of ups (e.g. new job, unexpected savings) and downs (e.g. job loss, unexpected bill). A technology-enhanced treatment model that is more responsive to variability in those emergent stressors (e.g. Guan, Park, & Chorpita, 2019; Lind et al., 2020), as well as its impact on day-to-day parenting and skill use, is likely critical to achieving and preserving treatment gains.

Acknowledgments

This study was funded by the National Institute of Mental Health (NIMH) R01MH100377 (D.J.J.; ClinicalTrials.gov Identifier: NCT02191956). Other support was provided by NIMH R21MH113887 (D.J.J.; ClinicalTrials.gov Identifier: NCT03597789), the National Institute of Child Health and Human Development (R.L.; T32HD007377), the National Science Foundation (A.H.; DGE-1650116), the National Institute of Minority Health and Health Disparities (J.P., PI; R01MD015401), and the National Institute on Drug Abuse (M.H.; T32DA043449). The authors are grateful to the families, assessors, and therapists who participated in this project for their time and valuable contributions. The authors have declared that they have no competing or potential conflicts of interest.

Correspondence

Justin Parent, Warren Alpert Medical School, Brown University, 1 Hoppin St #204, Providence, RI, USA; Email: justin_parent@brown.edu

Key points

- Behavioral parent training (BPT) is the standard of care for early onset disruptive behavior disorders, but waning treatment effects remain a challenge.
- An RCT compared a standard BPT program, Helping the Noncompliant Child (HNC), to Technology-Enhanced HNC (TE-HNC) in a clinical sample on parent proficiency of program skills and child behavior at post-treatment and 3- and 6-month follow-up.
- Both programs yielded improvements in parent skill proficiency and child problem at post-treatment, but TE-HNC families evidenced greater maintenance of some treatment gains at follow-up.
- The findings suggest that a technology-enhanced treatment model may allow for deeper learning of parent skills and improvement in child behavior.

References

- Anton, M.T., & Jones, D.J. (2019). Parent-therapist alliance and technology use in behavioral parent training: A brief report. *Psychological Services, 16*, 260–265.
- Blumberg, S.J., & Luke, J.V. (2018). *Wireless substitution: Early release of estimates from the National Health Interview Survey, January–June 2018*. US Department of Health and Human Services, Centers for Disease Control and Prevention. Available from: <https://www.cdc.gov/nchs/data/nhis/earlyrelease/wireless201812.pdf>
- Chacko, A., Isham, A., Cleek, A.F., & McKay, M.M. (2016). Using mobile health technology to improve behavioral skill implementation through homework in evidence-based parenting intervention for disruptive behavior disorders in youth: Study protocol for intervention development and evaluation. *Pilot Feasibility Studies, 2*.
- Chacko, A., Wymbs, B.T., Chimiklis, A., Wymbs, F.A., & Pelham, W.E. (2012). Evaluating a comprehensive strategy to improve engagement to group-based behavioral parent training for high-risk families of children with ADHD. *Journal of Abnormal Child Psychology, 40*, 1351–1362.
- Chorpita, B.F., Daleiden, E.L., Ebesutani, C., Young, J., Becker, K.D., Nakamura, B.J., ... & Starace, N. (2011). Evidence-based treatments for children and adolescents: An updated review of indicators of efficacy and effectiveness. *Clinical Psychology: Science and Practice, 18*, 154–172.
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* (2nd edn). Hillsdale, NJ: Lawrence Erlbaum.
- Cohen, M.A., & Piquero, A.R. (2009). New evidence on the monetary value of saving a high risk youth. *Journal of Quantitative Criminology, 25*, 25–49.
- Comer, J.S., Chow, C., Chan, P.T., Cooper-Vince, C., & Wilson, L.A. (2013). Psychosocial treatment efficacy for disruptive behavior problems in very young children: a meta-analytic examination. *Journal of the American Academy of Child and Adolescent Psychiatry, 52*, 26–36.
- Cui, Z., & Gong, G. (2018). The effect of machine learning regression algorithms and sample size on individualized behavioral prediction with functional connectivity features. *NeuroImage, 178*, 622–637.
- de Graaf, I., Speetjens, P., Smit, F., de Wolff, M., & Tavecchio, L. (2008). Effectiveness of the Triple P Positive Parenting Program on behavioral problems in children: A meta-analysis. *Behavior Modification, 32*, 714–735.
- Eyberg, S.M., & Pincus, D. (1999). *Eyberg child behavior inventory and Sutter-Eyberg Behavior Inventory-Revised: Professional manual*. Odessa, FL: Psychological Assessment Resources.
- Eyberg, S.M., & Robinson, E.A. (1983). Conduct problem behavior: Standardization of a behavioral rating scale with adolescents. *Journal of Clinical Child Psychology, 12*, 347–354.
- Fergusson, D.M., Horwood, L.J., & Ridder, E.M. (2005). Show me the child at seven: The consequences of conduct problems in childhood for psychosocial functioning in adulthood. *Journal of Child Psychology and Psychiatry, 46*, 837–849.
- Funderburk, B.W., Eyberg, S.M., Rich, B.A., & Behar, L. (2003). Further psychometric evaluation of the Eyberg and Behar rating scales for parents and teachers of preschoolers. *Early Education and Development, 14*, 67–81.
- Gelman, A., & Rubin, D.B. (1992). Inference from iterative simulation using multiple sequences. *Statistical Science, 7*, 457–511.
- Georgeson, A.R., Highlander, A., Loiselle, R., Zachary, C., & Jones, D.J. (2020). Engagement in technology-enhanced interventions for children and adolescents: Current status and recommendations for moving forward. *Clinical Psychology Review, 78*, 101858.
- Guan, K., Park, A.L., & Chorpita, B.F. (2019). Emergent life events during youth evidence-based treatment: Impact on future provider adherence and clinical progress. *Journal of Clinical Child and Adolescent Psychology, 8*, S202–S214.
- Jacobson, N.S., & Truax, P. (1991). Clinical significance: A statistical approach to defining meaningful change in psychotherapy research. *Journal of Consulting and Clinical Psychology, 59*, 12–19.
- Jones, D.J., Forehand, R., Cuellar, J., Kincaid, C., Parent, J., Fenton, N., & Goodrum, N. (2013). Harnessing innovative technologies to advance children's mental health: Behavioral parent training as an example. *Clinical Psychology Review, 33*, 241–252.
- Jones, D.J., Forehand, R., Cuellar, J., Parent, J., Honeycutt, A., Khavjou, O., ... & Newey, G.A. (2014). Technology-enhanced program for child disruptive behavior disorders: Development and pilot randomized control trial. *Journal of Clinical Child and Adolescent Psychology, 43*, 88–101.
- Jones, D.J., Loiselle, R., Zachary, C., Georgeson, A.R., Highlander, A., Turner, P., ... & Forehand, R. (2021). Optimizing engagement in behavioral parent training: Progress toward a technology-enhanced treatment model. *Behavior Therapy, 52*, 508–521.
- Kaminski, J.W., & Claussen, A.H. (2017). Evidence base update for psychosocial treatments for disruptive behaviors in children. *Journal of Clinical Child & Adolescent Psychology, 46*, 477–499.
- Keller, B.T., & Enders, C.K. (2019). *Blimp user's guide (version 2.0)*. Los Angeles.

- Khavjou, O., Forehand, R., Loiselle, R., Turner, P., Buell, N., & Jones, D.J. (2020). Helping the noncompliant child: An updated assessment of program costs and cost-effectiveness. *Children and Youth Services Review, 114*, 105050.
- Khavjou, O.A., Turner, P., & Jones, D.J. (2018). Cost effectiveness of strategies for recruiting low-income families for behavioral parent training. *Journal of Child and Family Studies, 27*, 1950–1956.
- Lavigne, J.V., Lebailly, S.A., Gouze, K.R., Binns, H.J., Keller, J., & Pate, L. (2010). Predictors and correlates of completing behavioral parent training for the treatment of oppositional defiant disorder in pediatric primary care. *Behavior Therapy, 41*, 198–211.
- Leijten, P., Raaijmakers, M.A., de Castro, B.O., & Matthys, W. (2013). Does socioeconomic status matter? A meta-analysis on parent training effectiveness for disruptive child behavior. *Journal of Clinical Child and Adolescent Psychology, 42*, 384–392.
- Lind, T., Lau, A.S., Gomez, C., Rodriguez, A., Guan, K., Chlebowsky, C., ... & Brookman-Frazee, L. (2020). Emergent life events in the delivery of a caregiver-mediated evidence-based intervention for children with autism spectrum disorder in publicly funded mental health services. *Autism, 24*, 1286–1299.
- Lindhiem, O., Bennett, C.B., Rosen, D., & Silk, J. (2015). Mobile technology boosts the effectiveness of psychotherapy and behavioral interventions: A meta-analysis. *Behavior Modification, 39*, 785–804.
- Lundahl, B., Risser, H.J., & Lovejoy, M.C. (2006). A meta-analysis of parent training: Moderators and follow-up effects. *Clinical Psychology Review, 26*, 86–104.
- Masarik, A.S., & Conger, R.D. (2017). Stress and child development: A review of the family stress model. *Current Opinion in Psychology, 13*, 85–90.
- McCabe, K.M., Yeh, M., & Zerr, A.A. (2020). Personalizing behavioral parent training interventions to improve treatment engagement and outcomes for culturally diverse families. *Psychology Research and Behavior Management, 13*, 41–53.
- McMahon, R.J., & Forehand, R.L. (2003). *Helping the noncompliant child: Family-based treatment for oppositional behavior* (2nd edn). New York: Guilford Press.
- Merikangas, K.R., Nakamura, E.F., & Kessler, R.C. (2009). Epidemiology of mental disorders in children and adolescents. *Dialogues in Clinical Neuroscience, 1*, 7–20.
- Muthén, L.K., & Muthén, B.O. (2017). *Mplus: Statistical analysis with latent variables: User's guide (Version 8)*. Los Angeles: Authors.
- Nixon, R.D.V., Sweeney, L., Erickson, D.B., & Touyz, S.W. (2003). Parent-child interaction therapy: A comparison of standard and abbreviated treatments for oppositional defiant preschoolers. *Journal of Consulting and Clinical Psychology, 71*, 251–260.
- Ogders, C.L., Caspi, A., Broadbent, J.M., Dickson, N., Hancox, R.J., Harrington, H.L., ... & Moffitt, T.E. (2007). Prediction of differential adult health burden by conduct problem subtypes in males. *Archives of General Psychiatry, 64*, 476–484.
- Ortiz, C., Vidair, H.B., Acri, M., Chacko, A., & Kobak, K. (2020). Pilot study of an online parent-training course for disruptive behavior with live remote coaching for practitioners. *Professional Psychology: Research and Practice, 51*, 125–133.
- Pew Research Center (2015). *The smartphone difference*. Available from: <http://www.pewinternet.org/2015/04/01/us-smartphone-use-in-2015/>
- Pew Research Center (2019). *Mobile fact sheet*. Available from: <http://www.pewinternet.org/fact-sheet/mobile/>
- Piquero, A.R., Jennings, W.G., Diamond, B., Farrington, D.P., Tremblay, R.E., Welsh, B.C., & Gonzalez, J.M.R. (2016). A meta-analysis update on the effects of early family/parent training programs on antisocial behavior and delinquency. *Journal of Experimental Criminology, 12*, 229–248.
- Polanczyk, G.V., Salum, G.A., Sugaya, L.S., Caye, A., & Rohde, L.A. (2015). Annual research review: A meta-analysis of the worldwide prevalence of mental disorders in children and adolescents. *Journal of Child Psychology and Psychiatry, 56*, 345–365.
- Preacher, K.J., Wichman, A.L., MacCallum, R.C., & Briggs, N.E. (2008). *Latent growth curve modeling* (No. 157). New York: Sage.
- Shaw, D.S., & Shelleby, E.C. (2014). Early-starting conduct problems: Intersection of conduct problems and poverty. *Annual Review of Clinical Psychology, 10*, 503–528.
- Timmons, A.C., Han, S.C., Chaspari, T., Kim, Y., Pettit, C., Narayanan, S., & Margolin, G. (2019). Family-of-origin aggression, dating aggression, and physiological stress reactivity in daily life. *Physiology & Behavior, 206*, 85–92.
- Webster-Stratton, C., & Hammond, M. (1997). Treating children with early-onset conduct problems: A comparison of child and parent training interventions. *Journal of Consulting and Clinical Psychology, 65*, 93–109.
- Williams, G.C., Lynch, M., & Glasgow, R.E. (2007). Computer-assisted intervention improves patient-centered diabetes care by increasing autonomy support. *Health Psychology, 26*, 728–734.

Accepted for publication: 10 November 2021